

Original Article

Zinc Impurity in Drinking Water and Its Toxic Effect on Human Health

Mahipal Singh Sankhla^{1*}, Rajeev Kumar² and Lalit Prasad³

¹Research Scholar, ²Associate Professor, Division of Forensic Science, SBAS, Galgotias University, Greater Noida, U.P., India

³Associate Professor, Division of Chemistry, SBAS, Galgotias University, Greater Noida, U.P., India

*Corresponding author email id: mahipal4n6@gmail.com

Received: 21-09-2019; Accepted: 26-09-2019

ABSTRACT

The levels of zinc (Zn) pollution in several water sources as ground, river, tap, lake water etc. Zinc are possibly toxic and are moved to the nearby environment through different pathways. Among mining activity, waste dump contaminants one of the most hazardous groups for of their determined nature, tendency, toxicity, to collect in creatures and undertake food chain increase and more still, they are non-degradable. Zinc concentration are found in water are higher than the permissible limits of WHO, USEPA that gives an indication of hazardous risk to human health. This review study also reveals that zinc contamination through water and major health effects on human health.

Keywords: Heavy metals, Zinc, Environment, Water, Toxicity

INTRODUCTION

Water is a valuable and most frequently used resource. Zinc is one of the most abundant compounds on earth, and covers two thirds of the earth's surface ^[1]. Humans lack access to clean and pure safe water ^[2-3]. Zinc is a very communal pollutant in environment; its occurrence may be impends the water ecological environment. Consequently, much study effort has been directed toward the spreading of Zn in water environment. Anthropogenic actions counting municipal wastewater releases, coal-burning power plants; industrial methods involving metals; and atmospheric outcome are the main source of Zn contamination ^[4]. Extreme discharge of zinc contaminate the surface water and subsurface environment and contribute to groundwater pollution. Groundwater is often extremely polluted near mines of sulfide minerals ^[5]. Development is the growth of rational and financial trade

that changes a people assembly from a rural culture into a current one. India began its manufacturing development after independence. The manufacturing strategy resolve of 1948 obvious the opening of the progress of manufacturing policy ^[6]. Mining Auctions, water bodies are most insistently polluted. The metals are taken as either dissolved species in water or as an essential part of on hold sediments. They may then be kept in bed residues or seep into the groundwater, mainly wells and springs; and the amount of pollution will be gritty by on the nearness of the well and spring to the mining site. As an outcome of these issues, metal absorption in the natural environment changes in space and time. In fact, during the last few decades, industrial and urban actions have contributed to the growth of metals pollution into semi-arid environment and have directly partial urban ecosystems producing toxic, mutagenic or carcinogenic effects to the human health dependent on the elements

properties. The probable effect of these elements may be showed by risk assessment ^[7]. The main sources of anthropogenic action zinc are found in the environment from metal smelters and mining actions ^[8]. In natural waters, zinc can be found in some chemical methods, such as hydrated ions, metal-inorganic complexes, or metal-organic complexes. Hydrated zinc cations may be hydrolyzed to form zinc hydroxide or zinc oxide. In anaerobic environments, Zinc sulfide may be formed ^[9].

Major sources of Water Contamination

Sewage and wastewater

Regular local households and outdated rural practices produce wastewater proficient of contaminating rivers tributaries and lakes. In the lack of fresh water and hygienic situations, efficient elimination of sewage is non-existing, and encourages water-borne ailments ^[10].

INDUSTRIAL WASTE

Industrial wastes are the main sources which pollute water. Presenting extremely toxic contaminants affecting humans and the environment ^[11]. The industrial use of freshwater in manufacturing and treating and through the contaminated water into lakes, rivers and groundwater ^[12].

MINING ACTIVITIES

Zinc is found in physical background on earth's crust and can consequently enter water sources through natural procedures and the example of sometimes heavy rains and water can leach zinc out of natural sources ^[13]. The methods are improved when this physical condition of earth is troubled by financial actions such as mining. These procedures of description the mined-out range to water and air, and can lead to significances such as acid mine drainage. Low pH situations related with acid mine drainage mobilize heavy metals, containing radio nuclides where these are present in the mines ^[14].

Zinc Toxic Effect on Human Health

Gastrointestinal Effects

Various research have recommended that zinc inhalation

and drinking polluted water may origin signs of gastrointestinal diseases or modifications in gastrointestinal soft tissue. One example of A person who consumed around 3 grains of a zinc chloride defined critical signs that happened nearly instantly after interaction with the complex, containing scorching, mouth and esophagus was vomiting and paining ^[15].

Immunological and Lymph reticular Effects

Zinc play a major part the regular growth and preservation of the immune structure, such as in the lymphocyte reaction to mitogens and as a cofactor for the thymic hormone thymulin ^[16-17]. Metal smoke infection is assumed to be a resistant reaction to zinc oxide. A connection among the amount of floating zinc and quantity of all forms of T cells in the Broncho alveolar lavage liquid of people, probably associated to the start of metallic smoke infection, was seen in a critical period breathing study ^[18]. Decreased resistant reaction in people has been informed in a middle period oral study^[19].

Cardiovascular Effects

Reported signs in humans showing excessive concentration of zinc containing premature atrial strokes, hypertension subordinate to intravascular volume, hypovolemic shock wave (pulse over 120 beats per minute) and hypertension ^[20].

Carcinogenic Effect

The molecular mechanisms damages of zinc. DNA damage considered to defend against effects genetic constancy and purpose, improves the vulnerability to DNA-damaging agents and furthermore moves cellular differentiation, proliferation and apoptosis. This has led people to hazard that zinc deficiency is a probable risk factor for cancer ^[20].

Neurotoxicity

High level of Consumption zinc have caused in tiredness, dizziness, shocking, trouble in writing properly, anxiety, sadness, somnolence and dragging ^[20]. Confounding gait and illusions were informed in a specific who purposely breathe in metallic paint vaporizers ^[21]. The instantaneous

contact to copper and hydrocarbons. Common Signs of neurological toxicity (dizziness, pain, headache and tiredness) have been described by human's subsequent critical oral intake to zinc [22, 23]. Actual Restricted Information Recommend that overdoses oral intake of zinc can result in light neuron collapse and change of emission of the hypothalamus in rats [24, 25]. Further researches by all tracks of contact would be beneficial to decide if zinc contact to complexes would result in neurological toxicity [26].

Toxicokinetics

There is partial data on the toxic kinetic of zinc subsequent breathing or skin exposure. Level of zinc increases in urine or blood of people and the soft muscle of animals later drink contaminated water, breathing and skin contact to zinc, individually, show that zinc is engrossed by these paths. The zinc consume been broadly ingested toxicokinetic properties. The involvement of zinc after the abdominal area is homeo statically controlled; below regular physical situations, 20–30% of consumed zinc is absorbed. After the Zinc taken abdominal includes submissive dispersion and a carrier facilitated procedure. A amount of features impact the involvement of zinc contain soluble in the zinc complex as Interception such as phosphorus, calcium and nutritional phytates or fiber and accompaniments, such as picolinic acid, amino acids and prostaglandin E2. Absorption of the zinc is broadly dispersed all over the body. Zinc contented is leading in muscular, bones, abdominal tract, lung, brain, pancreas, kidney, and skin and cardiovascular. In plasma, two-thirds of the zinc is certain to albumin which signifies the metabolic activity group of zinc. This group of plasma zinc is generally declared to as lightly certain zinc since albumin has the capability to provide to certain zinc to muscles [26].

DISCUSSION

The water is the main route and source to intake the zinc directly. If they found zinc in excessive amount of the permissible levels according to procedures of drinking water given by WHO, USEPA. Several deaths due to high concentration of zinc toxicity to human. There is a

need of solid waste management in major human settlements so that dangerous chemicals are not pollute the water bodies. The type of contamination is dependent on different type of usage of water and other activities like Industrial, Agricultural and sewage. Several types of biological, organic, and inorganic contamination have major effects on water quality. Many cost-effectiveness are the important factors that play major roles in the assortment of the most appropriate treatment system for wastewater.

CONCLUSION

Zinc has naturally originated either from geological and chemical variance or tailings site. Zinc indicate an anthropogenic source due to the sulfide minerals oxidation in mine waste disposal site. Over the past year's environmental principles have developed more stringent, necessitating an enhanced better quality of removing toxic waste for treatment and zinc material to defense of the environment, human health and aquatic life. Optional that, mining wastes must be treated before discarding them to the land and natural water resources. This will protect human health along with the environment.

REFERENCES

- [1] Ramesh K, Elango, L. Impact of groundwater quality from industrial East Coastal Town, Southern India. *International Journal of Engineering Research and Applications*. 2014; 4(1): 346–354.
- [2] Bresline E. Sustainable water supply in developing countries. *Geological Society of America*. 201; 1(1): 194. <https://doi.org/10.1186/s40064-015-0867-0>
- [3] The National Academy of Science NAS (2009) National Academy of Science: Overview-Safe Drinking Water is Essential. The National Academies Press, Washington, D.C
- [4] Pertsemli E, Voutsas D. Distribution of heavy metals in Lake Doirani and Kerkini, Northern Greece. *Journal of Hazardous Materials*. 2014; 148(1): 529–37. <https://doi.org/10.1016/j.jhazmat.2007.03.019>
- [5] Ghadimi M, Ghomi AH. Identification of groundwater contamination sources of Lakan lead and zinc mine, Khomein Iran. 2012; 3(2): 121-134. doi: 10.22044/JME.2013.96

- [6] Singh J, Yadav H, Smarandache F. District level analysis of urbanization from rural-to-urban migration in the Rajasthan state. *Smarandache Notions Journal*. 2009; 1(2): 1–12.
- [7] Batayneh AT. Toxic (aluminum, beryllium, boron, chromium and zinc) in groundwater: health risk assessment. *International Journal of Environmental Science and Technology*. 2012; 9(1): 153–162. doi: 10.1007/s13762-011-0009-3.
- [8] ATSDR (Agency for Toxic Substances and Disease Registry) (1995). Toxicological profile for zinc. Public Health Service, U.S. Department of Health and Human Services, Atlanta, GA.
- [9] Callahan MA, Slimak MW, Gabel NW, May IP, Fowler CF, Freed JR 1979. Water-related environmental fate of 129 priority pollutants. Volume 1: Introduction and technical background, metals and inorganics, pesticides and PCBs [Final Report].
- [10] Beuchat LR. Vectors and conditions for preharvest contamination of fruits and vegetables with pathogens capable of causing enteric diseases. *British Food Journal*. 2006; 108(1): 38–53. doi 10.1108/00070700610637625
- [11] Jain AK, Gupta VK, Bhatnagar A. Utilization of industrial waste products as adsorbents for the removal of dyes. *Journal of Hazardous Materials*. 2003; 101(1): 31–42. [https://doi.org/10.1016/S0304-3894\(03\)00146-8](https://doi.org/10.1016/S0304-3894(03)00146-8)
- [12] Hines NW. Controlling industrial water pollution: color the problem green. *BCL Reviews*. 1967; 9(1): 553–6.
- [13] Sankhla MS, Kumari M, Nandan M, Kumar R, Agrawal P. Heavy Metals Contamination in Water and their Hazardous Effect on Human Health-A Review. *International Journal of Current Microbiology Applied Science*. 2016; 5(10): 759–766. doi: <http://dx.doi.org/10.20546/ijemas.2016.510.082>
- [14] Bagul VR, Shinde DN, Chavan RP, Patil CL, Pawar RK. New perspective on heavy metal pollution of water. *Journal of Chemical and Pharmaceutical Research*. 2015; 7(12): 700–705.
- [15] Chobanian SJ. Accidental ingestion of liquid zinc chloride: Local and Systemic Effects. *Annals of Emergency Medicine*. 1981; 10(1): 91–93. doi: [https://doi.org/10.1016/S0196-0644\(81\)80345-9](https://doi.org/10.1016/S0196-0644(81)80345-9)
- [16] Delafuente JC. Nutrients and immune responses. *Rheumatic Diseases Clinics of North America*. 1991; 17(2): 203–212.
- [17] Fraker PJ, Gershwin ME, Good RA, Prasad A. Interrelationships between zinc and immune function. In *Federation Proceedings*. 1986; 45(5): 1474–1479.
- [18] Blanc P, Wong H, Bernstein MS, *et al.* An experimental human model of metal fume fever. *Annals of Internal Medicine*. 1991; 114(1): 930–936. doi: 10.7326/0003-4819-114-11-930
- [19] Chandra RK. Excessive intake of zinc impairs immune responses. *JAMA* 1984; 252(1):1443–1446. doi:10.1001/jama.1984.03350110043027
- [20] Nriagu, J. Zinc toxicity in humans. School of Public Health, University of Michigan. 2007.
- [21] Wilde C. Aerosol metallic paints: Deliberate inhalation: A study of inhalation and/or ingestion of copper and zinc particles. *International Journal of Addiction*. 1975; 10(2): 127–134. <https://doi.org/10.3109/10826087509026709>
- [22] Murphy JV. Intoxication following ingestion of elemental zinc. *JAMA*. 1970; 212: 2119–2120. doi:10.1001/jama.1970.03170250073021
- [23] Potter JL. Acute zinc chloride ingestion in a young child. *Annals of Emergency Medicine*. 1981; 10(5): 267–269. doi: [https://doi.org/10.1016/S0196-0644\(81\)80055-8](https://doi.org/10.1016/S0196-0644(81)80055-8)
- [24] Kozik MB, Gramza G, Pietrzak M. Neurosecretion of the hypothalamo-hypophyseal system after intragastric administration of zinc oxide. *Folia histochemica et Cytochemica*. 1981;19(2): 115–22.
- [25] Kozik MB, Maziarz L, Godlewski A. Morphological and histochemical changes occurring in the brain of rats fed large doses of zinc oxide. *Folia Histochemica et Cytochemica*. 1980; 18(3): 201–6.
- [26] ATSDR (Agency for Toxic Substances and Disease Registry) (2005). Toxicological profile for zinc. Public Health Service, U.S. Department of Health and Human Services, Atlanta, GA.

How to cite this article: Mahipal Singh Sankhla, Rajeev Kumar and Lalit Prasad. Zinc Impurity in Drinking Water and Its Toxic Effect on Human Health. *Indian Internet Journal of Forensic Medicine & Toxicology*. 2019; 17(4): 84-87.