

INCLUSION OF CIDR IN OVSYNCH PROTOCOL TO IMPROVE FERTILITY IN POSTPARTUM SUBESTRUS BUFFALOES

K. RAVIKUMAR¹, S.A. ASOKAN² and C. VEERAPANDIAN³

Department of Animal Reproduction, Gynaecology and Obstetrics,
Madras Veterinary College, Chennai-600 007.

ABSTRACT

Twenty four postpartum subestrus buffaloes not showing observed estrus within 60 days after calving were randomly and equally divided into ovsynch-subestrus and ovsynch plus CIDR-subestrus groups. Ovsynch-subestrus group buffaloes were subjected to ovsynch protocol which consisted of 100 µg i.m. injection of GnRH on the day of start of synchronization (d 0), 25 mg i.m. injection of PGF_{2α} seven days later (d 7), another 100 µg i.m. injection of GnRH 48 h after PGF_{2α} (d 9) and timed insemination 16 to 18 h after second GnRH injection (d 10). Ovsynch plus CIDR-subestrus buffaloes were subjected to ovsynch plus CIDR protocol which included intra vaginal placement of CIDR device for a period of 7 days starting from first GnRH injection (d 0) to PGF_{2α} injection (d 7) for synchronization of ovulation. Blood samples were collected from all experimental animals on d -10 and d 0, d 7, and d 9 days after insemination for the estimation of progesterone. During the treatment protocol none of the buffaloes exhibited visible premature estrus signs. However progesterone estimation showed that 25 per cent had exhibited premature estrus. The estrus detection rates were 41.66 and 58.33 per cent in ovsynch-subestrus and ovsynch plus CIDR-subestrus respectively. Most buffaloes showed intermediate signs of estrus. The ovulatory response and first service conception rates were 83.33 and 33.33 and 100 and 41.66 per cent in ovsynch-subestrus and ovsynch plus CIDR-subestrus groups respectively.

Key words: Ovsynch, CIDR, Subestrus buffaloes, Fertility.

INTRODUCTION

The major limitation of GnRH and PGF_{2α} based synchronization protocols was the inability of GnRH to turnover dominant follicles late in the estrous cycle leading to premature estrus in 8 to 10 per cent of treated cows (Geary *et al.*, 2000). Alternatively the inclusion of an exogenous such as CIDR during the interval between GnRH and PGF_{2α} injection prevented premature estrus and increased estrus response and conception rates (Steckler *et al.*, 2002). Although much work has been done using ovsynch and ovsynch plus CIDR in synchronization of ovulation in cattle, information on their use in buffaloes especially postpartum lactating

subestrus Murrah buffaloes is limited. Hence the present study was taken up (i) To study the effect of ovsynch protocol in induction of ovulation in postpartum subestrus buffaloes. (ii) To study the effect of CIDR-B on estrus synchronization in ovsynch treated postpartum subestrus buffaloes. (iii) To compare fertility rate among ovsynch and ovsynch plus CIDR-B treated postpartum subestrus buffaloes.

MATERIALS AND METHODS

Twenty four apparently healthy lactating buffaloes maintained at Central Cattle Breeding Farm, Alamathi, with a history of absence of observed estrus signs within 60 days were selected and subjected to rectal examination twice at an interval of ten days. All the selected animals had corpus luteum during rectal examination. Similarly serum progesterone levels were estimated twice (d- 10 and d 0) to confirm cyclic activity of the animals. The selected postpartum subestrus

Part of M.V.Sc thesis of first author submitted to Tamilnadu Veterinary and Animal Sciences University.
Asst. Prof. Dept. of Clinics, VC&RI, Namakkal-637 001.
Professor,² Professor and Head, Dept. of ARGO, Madras Veterinary College, Chennai-600 007.

buffaloes were randomly and equally divided into ovsynch-subestrus and ovsynch plus CIDR-subestrus groups.

The ovsynch subestrus group buffaloes were injected to administration of 100 µg i.m. injection of GnRH (Gonadorelin Acetate, Fertilineâ, Vetoquinol, Canada) on the day of start of synchronization (d 0), 25 mg i.m. injection of PGF_{2α} (Dinoprost. tromethamin, Lytalyseâ, Pharmacia, Belgium) seven days later (d 7), another 100 µg i.m. injection of GnRH 48 h after PGF_{2α} (d 9) and timed insemination 16 to 18 h after second GnRH injection (d 10). The ovsynch plus CIDR group buffaloes were treated with same ovsynch protocol and in addition intra vaginal placement of CIDR-B device (EAZI-BREED CIDR™, Inter Ag, Hamilton, New Zealand) for a period of 7 days starting from 1 st GnRH injection (d 0) to PGF_{2α} injection (d 7). The experimental buffaloes were observed frequently for estrus signs during the period of treatment and after PGF_{2α} injection. The percentage of estrus detection rate was estimated as the number of buffaloes detected in estrus during the 66 h after PGF_{2α} administration. Onset of estrus was calculated in hours from the time of PGF_{2α} administration or / and CIDR removal to the time of first appearance of estrus signs. The intensity of estrus was studied using behavioral changes, physiological changes and gynaecological observations and it was scored as described by Rao and Rao (1981) with slight modifications.

Blood samples were collected from all experimental animals on d- 10, d 0, d 7, and d 9 and the sera samples were subjected for the estimation of progesterone. All the animals had more than 1 ng/ml progesterone either at d- 10 or d 0 indicating that they were in cyclicity without observable estrus signs. The serum progesterone level was estimated using solid-phase Radio Immuno Assay technique with the help of progesterone kits (Coat-A-Count, Diagnostic Products Corporation, USA). Ovulatory response was assessed by rectal examination at 10 days after induced ovulation which was later confirmed by serum progesterone levels. First service conception rate was calculated as percentage of animals that conceived to fixed time insemination at induced estrus in each group. Pregnancy

was confirmed by palpation per rectum at 60 days post insemination.

RESULTS AND DISCUSSION

In the present study, none of the animals exhibited visible premature estrus signs between d 0 and d 7 of the treatment period. However, Gabor *et al* (2002) reported that the incidence of premature estrus to be 11 to 14 per cent while using ovsynch protocols. Although in the present study no visible premature estrus signs were observed, progesterone levels on day 7 and day 9 indicated that 25.00 per cent of the ovsynch subestrus group animals had less than 1 ng/ml indicating that they exhibited premature estrus however none of the CIDR treated exhibited premature estrus. The inclusion of CIDR in the ovsynch protocol in the present study may have potentially prevented the occurrence of premature estrus in ovsynch plus CIDR buffaloes. Estrus detection rate, onset of estrus, duration of estrus and intensity of estrus are presented in table. An increased percentage of estrus detection rate of 16.67 per cent (table-I) for ovsynch plus CIDR-subestrus over the ovsynch-subestrus buffaloes might be due to the inclusion of CIDR in ovsynch protocol in the present study.

There was no significant difference with regards to onset of estrus among the two treatment groups. Stevenson *et al.* (1999) reported the interval to onset of estrus after PGF_{2α} was 54 ± 13 and 55 ± 4.4 h in ovsynch 33 (2nd GnRH 33 h after PGF_{2α}) and ovsynch 48 (2nd GnRH 48 h after PGF_{2α}) treated Holstein cows. The intensity of estrus was almost the same in ovsynch-subestrus and ovsynch plus CIDR-subestrus groups. Neglia *et al.* (2003) reported that mucus and estrus behaviour were only observed in rare occasions. However, 88 per cent of Italian Mediterranean buffaloes treated with ovsynch protocol had a tonic uterus on the day of AI.

The ovulatory response and first service conception rate were presented in table-I. The ovulatory response for ovsynch-subestrus buffaloes (83.33%) in the present study was similar to those reported by Steckler *et al.* (2002) in cows. Higher ovulatory responses have been reported by Berber *et al.* (2002)

(93.3 per cent) in buffaloes, while a lower ovulatory response (52.9 per cent) have been reported by Santos *et al.* (2002) in Holstein lactating cows. The ovulatory response of 100 per cent observed for ovsynch plus CIDR-subestrus buffaloes in the present study was higher than those reported by Steckler *et al.* (2002) (85 per cent) for cyclic postpartum cows. Between ovsynch and ovsynch plus CIDR groups a higher ovulatory response was observed for the ovsynch plus-CIDR group. The inclusion of progesterone in an ovsynch

protocol increased LH pulse frequency (Garcia-Winder *et al.*, 1986) thereby providing a favourable condition for the follicles to grow. As follicles grew, the increased production of estrogens (Garcia-Winder *et al.*, 1987) most likely increased the positive feed back of estradiol – 17 b on LH release from the pituitary once GnRH was injected on d 9. As a result of the increased concentrations of estradiol – 17 b in cows treated previously with progesterone, more LH was released after GnRH (Thompson *et al.*, 1999) which may have

TABLE : ESTRUS PATTERN, OVULATORY RESPONSE AND FIRST SERVICE CONCEPTION RATE IN OVSYNCH AND OVSYNCH PLUS CIDR TREATED SUBESTRUS BUFFALOES

Reproductive parameters	Estrus detection rates (per cent)	Onset of estrus		Intensity of estrus at the time of AI No. of animals (per cent)			Ovulatory response	First service conception rate
		Mean \pm SE (h)	Range (h)	Intense	Intermediate	Weak		
Ovsynch per cent (no./no.)	41.66 (5/12)	48.80 \pm 7.74	24-61	2 (16.67)	8 (66.66)	2 (16.67)	83.33 (10/12)	33.33 (4/12)
Ovsynch plus CIDR per cent (no./no.)	58.33 (7/12)	36.71 \pm 2.65	24-48	2 (16.67)	8 (66.66)	2 (16.67)	100 (12/12)	41.66 (5/12)

resulted in the increased ovulatory response. Higher conception rates following treatment with ovsynch have been reported by Berber *et al.* (2002) (61.7 per cent) in postpartum buffaloes and High conception rates of 49 and 72 per cent have been reported by Pursley *et al.* (2001) and Steckler *et al.* (2002) in ovsynch plus CIDR treated postpartum dairy cows as against the 41.66 per cent conception rate obtained in the present study for postpartum buffaloes.

The results of the present study have shown that the addition of progesterone in ovsynch plus CIDR protocol potentially could prevent the premature occurrence of estrus prior to or following PGF_{2a}. Ovsynch plus CIDR-subestrus group responded better with increased estrus response, higher ovulatory response and conception rates when compared to Ovsynch-subestrus group. Thus, to conclude, inclusion of a CIDR device with ovsynch protocol may be an ideal strategy for dealing with lactating postpartum subestrus buffaloes.

REFERENCES

- Berber, R.C.A., E.H.Madureira and Baruselli P.S. (2002). Comparison of two ovsynch protocols (GnRH versus LH) for fixed timed insemination in buffalo (*Bubalus bubalis*). *Theriogenology*, 57:1421 – 1430.
- Garcia – Winder, M., P.E.Lewis, D.R.Deaver, V.G.Smith, G.S.Lewis and Inskeep E.K. (1986). Endocrine profiles associated with lifespan and induced corpora lutea in postpartum beef cows. *J.Anim. Sci.*, 62 : 1353-1362.
- Garcia – Winder, M., P.E.Lewis, E.C.Townsend and Inskeep P E.K. (1987). Effects of norgestomet on follicular development in postpartum beef cows. *J.Anim. Sci.*, 64 : 1099-1109.
- Geary, T.W.,E.R.Downing, J.E.Bruehmet and Whitter J.C. (2000). Ovarian and estrus response of suckled beef cows to the select synch estrus synchronization protocol. *Prof.Anim. Sci.*, 16 : 1-5.
- Gabor, Gy., J.P.Kastelic, S.Pinter, F.Szasz, Edit Azigeti and Solymosi (2002). Improving reproductive performance

in lactating dairy cows by synchronising ovulation or inducing oestrus. *Acta Veterinaria Hungarica*, **50**(2) : 231-234.

- Neglia, G., B.Gasparini, R.Dpalo, C.DeRosa, L.Zicarelli and Campnile G. (2003). Comparison of pregnancy rates with two estrus synchronization protocols in Italian Mediterranean Buffalo Cows. *Theriogenology*, **60**: 125-133.
- Pursley, J.R., P.M. Fricke, H.A. Garverick, D.J. Kesler, J.S. Ottobre, J.S. Stevenson and M.C. Wiltbank (2001). NC-113 Regional Research Project. Improved fertility in anovulatory lactating dairy cows treated with exogenous progesterone during ovsynch. *J. Dairy Sci.* (Midwest Branch ADSA Meetings, Des Moines, IA, Abstract 251 p.63).
- Rao, S.V. and A.R. Rao (1981). Oestrus behaviour and ovarian activity of crossbred heifers. *Indian Vet. J.*, **58** : 881-884.
- Santos, R.M., J.L.M.Vasconcelos, E.P.B. C.Silva, M.Meneghetti, N.Ferreira Junior, Oliveira C.A. (2002). Evaluation of the efficiency of D-Cloprostenol or Cloprostenol sodium and different doses of gonadorelin (0.10 mg vs 0.25 mg) in the ovsynch protocol in Holstein cows. *Anim. Breed. Abstr.*, **70** (4) : 325.
- Steckler, T.L., T.F. Lock, G.C. Mcloy and Kesler D.J. (2002). Efficacy of ovsynch alone (or) in combination with intravaginal progesterone inserts in dairy cows. (Available at : <http://www.selectsires.com/selections/2002-q3> - Page 7-10.html). Accessed May 2003.
- Stevenson, J.S., Y.Kobayashi and Thompson K.E. (1999). Reproductive performance of dairy cows in various programmed breeding systems including ovsynch and combinations of gonadotropin - releasing hormone and prostaglandin F_{2α}. *J.Dairy Sci.*, **82** : 506 - 515.
- Thompson, K.E., J.S. Stevenson, G.C. Lamb, D.M. Grieger and Loest C.A. (1999). Follicular, hormonal and pregnancy responses of early postpartum suckled beef cows to GnRH, norgestomet and prostaglandin F_{2α}. *J. Anim. Sci.*, **77** : 1823-1832.

INSTRUCTIONS FOR CHAPTER SECRETARY / TREASURER

- The Chapter Secretary / Treasurer, receiving the membership fee of newly enrolled life members, may please send the name to Editor Immediately. The annual subscription fee for the journal may be remitted in full to us along with the names of the members, so that the journal is dispatched to them. The money may be send by draft or money transfer in favour of Editor IJAR payable at Chennai

Thanks

Sd/
Editor