

Prevalence of Malnutrition in *Bhal* Region of Gujarat State

Gayatree Jadeja¹ and M. S. Jakasaniya²

ABSTRACT

There has been increasing evidence that development of underweight and obesity has deleterious social and health consequences. The study was conducted to determine the prevalence of underweight and obesity in rural areas of Bhal region of Gujarat state. Total of 1213 subjects of four different age groups were selected from different 80 villages of rural area. The height, weight were measured for all age groups. Mid upper arm circumference, hair colour and skin texture were observed for preschool children as well as the blood pressure was measured for adults above 40 years. All parameters were correlated with the body mass index. The results found were alarming. Majority of girls had malnourishment symptoms compared to boys. More of tobacco addiction was found in adult male and 40 to 49 years adults were suffering from high blood pressure problem. Illiteracy was higher in female which is indirectly responsible for the health of the community.

Keywords: Malnutrition, body Mass Index

INTRODUCTION

Agricultural progress in the last decade has made India self-sufficient in major food grains. Yet undernutrition continues to be major nutritional problem especially in rural populations. High prevalence of low birth weight, high morbidity and mortality in children and poor maternal nutrition of the other continue to be major nutritional concerns in India. For India, published data show that increased mortality is associated with low BMI (NIN Annual Report, 1991) and increased incidence of low birth weight with increasing reduction of the BMI of pregnant women (Naidu and Rao, 1994). While we are in the midst of combating these long-standing problems of undernutrition in children and women, a new situation has arisen. The World health bank has predicted that coronary heart disease will become the leading cause of premature death in India by 2015 and that the maximum number of diabetic patients in the world will be in India (Bulatao and Stephens, 1992).

Gujarat has a high level of child malnutrition which is 47 per cent as compared to the national average 21 per cent (World Bank Report, 2005). In Gujarat, it is seen that malnutrition is not only affecting children but adults are also suffering from many macro and micro nutrient

deficiencies (NFHS III, 2005). The present study aimed in the light of the above issues and discuss strategies for achieving better health of our people.

METHODOLOGY

The prevalence of underweight and malnutrition is increasing rapidly, especially in underdeveloped countries. In describing the nutritional status of a population the body mass index (BMI) represents the most complete indicator and, at the same time, is the easiest to use. The study was conducted with the aims to find out the health status of all age group people by measurement of their height, weight, Body Mass Index and Blood pressure, to collect the family, medical and dietary history using personnel interview technique and to correlate the collected information with the health status of individual.

Sample Selection: Taking into consideration the time restraints and convenience, most of the villages from Bhal region Dholka taluka of Ahmedabad district were selected. Twelve hundred and thirteen subjects (1213) of different age groups were interviewed who were from different 80 villages of Bhal region. For the preschool and school children selection the schools were purposively selected and then the subjects were randomly selected.

¹SMS Home Science, ²Incharge Programme Coordinator, Krishi Vigyan Kendra, AAU, Arnej-382230 (Gujarat)

For the adolescents the purposive selection was done from the school as well as from the rural areas. Adults were selected those who were the participants of on and off campus training. The Height and weight were measured using standard techniques. Nutritional assessment was done on the basis of Body Mass Index (BMI). For the preschool children of 2 to 5 years total of 106 (53 boys and 53 girls) were selected from six different Anganwadies and their Mid upper Arm Circumference (MUAC), Head circumference, hair color and skin texture were observed. For school children 262 (136 boys and 126 girls) of 6 to 12 years and adolescents 222 (117 boys and 105 girls) of 13 to 17 years the subjects were selected from five different schools and their hair color and skin texture were visually observed. The adult group of 18 years and above subjects 623 (382 male and 241 female) were interviewed for the general questions like education, number of family members, suffering from any disease, any addiction (Tobacco/Mava/Masala etc). All these parameters were correlates with their BMI status. The blood pressure was also measured using the digital blood pressure measurement unit for the adult group.

Following standard techniques were used for measurements

Height: Height in centimeters was marked on a wall with the help of a measuring tape. All subjects were measured against the wall without foot wear and with heels together and their heads positioned so that the line of vision was perpendicular to the body. A metal scale was brought down to the topmost point on the head. The height was recorded to the nearest 1 cm.

Weight: The weight was measured using a weighing machine (Bathroom Scale) with an accuracy of ± 100 gm. The subjects were asked to remove their footwear before measuring their weight. The scales were recalibrated after each measurement. Accuracy of the weighing scale was verified from time to time against known weights

Body Mass Index (BMI): BMI of the study subject was calculated by using the formula $\text{weight (kg)} / \text{height}^2 (\text{m}^2)$. For grading proposed criteria of BMI of Asians and CDC (2010) was adopted. Children (6 to 17 years) and adults (18 and above years) with BMI below 18.5 were considered underweight whereas BMI at or above 25 were considered overweight.

Blood Pressure: Blood pressure of adults above 40 years (123 female and 248 male) were measured using the Fully Automatic Blood Pressure Monitoring instrument (Lloyd pharmacy, UK). For systolic and diastolic pressure grading proposed criteria of blood pressure cut off points

was adopted.

Hair Color and Skin texture: The visual observation was done for the hair color (Rusty red, light and dark bands of color) and skin texture (dry skin, wrinkled skin).

Statistical Analysis: Data thus generated were analysed using SPSS software.

RESULTS AND DISCUSSION

Group I (Pre-school children, 2-5 years): Malnutrition substantially raises the risk of infant and child deaths and increase the vulnerability to a variety of diseases in later life. Children who are undernourished and underweighted are likely to be less clever than if they are well fed (WHO report). Even though the problem of fetal growth may be taken care off, rural conditions are such that poor growth is a characteristic phenomenon among pre-school children and it may be interesting to examine postnatal effect of chronic undernutrition on adult size and health.

In this study, the pre-school children (53 girls and 53 boys) of six different anganwadies were examined for height, weight, hair color and skin texture. Table 1 data indicated that around 83 per cent girls and 84 per cent boys were underweight and 62 per cent girls and 67 per cent boys had not attained the normal height. This indicates that the early years of life are critical for rural children may miss the 'second opportunity' for catch-up growth during adolescence due to stunting in early life.

Mid upper arm circumference (MUAC) of girls was 5.6 per cent lower than the normal and 1.9 per cent in the pre-school boys which indicates that moderate malnutrition symptoms. Same way the head circumference shows that 33.9 per cent girls and 11.3 per cent boys were suffered from moderate to severe malnutrition. The symptoms found in girl child were higher than the boys.

Prolonged protein deficiency in the diet results in Kwashiorkor. This disease involves multiple symptoms including the reduction of pigment production and incorporation into hair fiber. Normally dark brown hair becomes a rusty red. Light colored hair become blonde. The flag sign sometimes seen in kwashiorkor involves alternating light and dark bands of color along individual hair fibers. The flag sign is associated with intermittent protein malnutrition. Presumably normal hair color is produced when protein intake is adequate. Severe vitamin B12 deficiency has been reported by some as a potential promoter of gray hair. Overall, it is extremely unusual to see hair color changes as a result of dietary deficiencies.

The results indicate that around 62 per cent girls and 37 per cent boys have brown rusty hair color and they were also of underweight and stunted height. So this may be an indication of protein malnutrition. The girl child has higher number of protein malnourishment compared to boys.

Table 1: Different anthropometric measurements of preschool children (2-5 years)

Anthropometric Parameters	Group I (Pre-school children)		
	Girl (n=53)	Boy (n=53)	Total (n=106)
Under weight	83.1 % (44)	84.9 % (45)	83.9 % (89)
Stunted (Lower height)	62.3 % (33)	67.9 % (36)	65.1 % (69)
Mid Upper Arm Circumference (MUAC) (< 12.5cm)	5.6 % (3)	1.9 % (1)	3.8 % (4)
Head Circumference (<45.5 cm)	33.9 % (18)	11.3 % (6)	22.6 % (24)
Brown Hair	62.3 % (33)	37.7 % (20)	50.0 % (53)
Dry Skin	67.9 % (36)	45.3 % (24)	56.6 % (60)

Figures in parentheses represent the number of subjects involved

Vitamin deficiency and dry skin can, quite literally, go hand in hand. The appropriate dosage of different vitamins and nutrients are essential for proper health and inadequate supplies of certain vitamins can result in dry skin. In our results we had observed that 67 per cent girls and 45 per cent boys were having the dry skin problem may because of vitamin deficiencies. Group II (School children, 6-12 years): School children (6 to 12 years) from different five schools were examining for anthropometric measurements.

The body mass index was calculated using height and weight measurements and the data were categorized for underweight, normal and overweight. Table 2 results indicate that out of 262 school children 97 per cent school children were in the underweight category. Almost all girls were of underweight category. So this is the indication of severe malnutrition problem. Brown rusty hair problem was observed in 57 per cent school children and the 61 per cent children were having the dry skin which was the indications of protein malnutrition and vitamin deficiencies.

When the body mass index was compared with the hair and skin observations the data indicate that around 97 per cent and 98 per cent school children were in the underweight category, respectively which may be the result of protein and vitamin deficiencies.

Table 2: Anthropometric measurements of school children (6-12 years)

Anthropometric measurements	Group II (School Children)		
	Girl (n=126)	Boy (n=136)	Total (n=262)
BMI			
< 18.5 (Under weight)	99.2 % (125)	96.4 % (131)	97.7 % (256)
18.5 to 24.9 (Normal)	0.8 % (1)	2.9 % (4)	1.9 % (5)
≥ 25 (Over weight)	0.0 % (0)	0.7 % (1)	0.4 % (1)
Brown hair	71.4 % (90)	44.8 % (61)	57.6 % (151)
Dry Skin	65.1 % (82)	57.4 % (78)	61.1 % (160)

Figures in parentheses represent the number of subjects involved

Group III (Adolescence, 13 to 17 years): Adolescence is known to be a “second opportunity” for growth as it facilitates catch-up growth for children experiencing nutrition deficits in their early life. However, as discussed above, stunting appeared to be a persistent phenomenon beyond the early life among rural children and it had significant impact at 10+ years of age. Thus the majority of children enter adolescence with poor nutritional status (Joshi *et al.* 1998, Rao *et al.* 1998a,b, 2000b; Kanade *et al.* 1999). We had therefore, examined the adolescence of 13 to 17 years (105 girls and 117 boys) from three different schools.

Table 3: Anthropometric measurements of adolescence (13-17 years)

Anthropometric measurements	Group III (Adolescence)		
	Girl (n=105)	Boy (n=117)	Total (n=222)
BMI			
< 18.5 (Under weight)	75.2 % (79)	83.8 % (98)	79.7 % (177)
18.5 to 24.9 (Normal)	23.8 % (25)	15.4 % (18)	19.4 % (43)
≥ 25 (Over weight)	0.95 % (1)	0.8 % (1)	0.9 % (2)
Brown hair	52.4 % (55)	31.6 % (37)	41.4 % (92)
Dry Skin	47.6 % (50)	19.7 % (23)	32.8 % (73)

Figures in parentheses represent the number of subjects involved

Table 3 indicates the body mass index of adolescence. 79 per cent adolescence was in the underweight category. Here the numbers were higher for boys than the girls. In adolescent girls, short stature that persists into adulthood is associated with increased risk of adverse reproductive outcomes (Thance *et al.*, 1997, Kirchengast and Winkler 1996).

Adolescence of 41 per cent were suffered from protein malnutrition as they had brown rusty hair as well as 32 per cent adolescence were having the vitamin deficiency because they have dry skin. Brown rusty hair and dry skin were observed more in the adolescent girls compared to adolescent boys. The adolescence that has

brown rusty hair (82.6) was having the underweight problem same way 84 per cent adolescence have dry skin may because of vitamin deficiencies. Adolescence boys were higher in underweight category compared to adolescence girls. Adolescence demands high level of activity and growth, dietary requirements both quantitatively as well as qualitatively are of great importance. A failure to consume an adequate diet during adolescence can potentially retard growth (Johnson et al., 1994).

Group IV (Adults, 18 years and above): The retrospective nature of observations based on which the fetal origins theory is proposed, overlooks the confounding effects of lifestyle and environmental factors in determining the risks for adult diseases. Continuous exposure to poor environmental conditions, morbidity and undernutrition in rural India, need investigation of postnatal influences in determining the risks associated with adult diseases. In this study 623 adults were involved out of them 242 were female and 382 were male subject. Their body mass index was calculated using the height and weight data and the other information like Blood pressure measurements of 40+adults, education, age, any addiction (mava/masala/tobacco etc.) were also collected. All the data regarding the other information were correlates with the BMI status of adults for getting the idea about the health status of adults. Table 4 indicates the body mass index data of adult category. 23 per cent adults were in underweight category whereas 21 per cent were in overweight category. The female subjects were higher in both the underweight and overweight category compared to male subjects.

Table 4: Body Mass Index of Adult group (≥18 years)

BMI	Group IV Adult ≥18 years		
	Female	Male	Total
<18.5 (Under weight)	29.8 % (72)	18.8 % (72)	23.2 % (144)
18.5 to 24.9 (Normal)	43.3 % (104)	62.3 % (238)	54.9 % (342)
≥ 25 (Over weight)	26.9 % (65)	18.9 % (72)	21.9 % (137)

Figures in parentheses represent the number of subjects involved

When the underweight and overweight subjects were age wise categorized we had observed that more of female in the age group of 18 to 39 years were of underweight subjects and more overweight female and male subjects were in 40 to 59 years age group. The 371 adults who are ≥ 40 years (123 female and 248 male) were examined for the blood pressure. Results show that 27 per cent adults had high systolic blood pressure and 31 per cent had high diastolic blood pressure. 70 per cent and 67 per cent adults were having the normal systolic and diastolic blood pressure respectively. In comparison of high blood

pressure with the different age groups the data revealed that 46 per cent to 49 per cent adults of 40 to 49 years were suffered from high systolic and diastolic blood pressure. Very few 60+ adults was having the high blood pressure.

Table 5: Blood Pressure prevalence in Adult (≥40 years)

Blood Pressure	Group IV Adult ≥ 40 years		
	Female (n= 123)	Male (n=248)	Total (n=371)
Systolic Blood Pressure			
≤ 90 (Lower side)	0.8 % (1)	2.5 % (6)	1.9 % (7)
91-129 (Normal Blood Pressure)	74.8 % (92)	68.1 % (169)	70.4 % (261)
≥ 130 (Higher side)	24.4 % (30)	29.4 % (73)	27.7 % (103)
Diastolic Blood Pressure			
≤ 60 (Lower side)	0.8 % (1)	1.6 % (4)	1.4 % (5)
61-84 (Normal Blood Pressure)	73.2 % (90)	64.1 % (159)	67.1 % (249)
≥ 84 (Higher side)	26.0 % (32)	34.3 % (85)	31.5 % (117)

Figures in parentheses represent the number of subjects involved

Adults with high blood pressure were compare with the body mass index and it indicated that more percentage of females were in the overweight category whereas the males were in the normal weight category. The information regarding the addiction of tobacco was also collected from adult male group. Results show that out of 382 adult male 212 *i. e.* 55 per cent were having the addiction of tobacco. When the data were categorized age wise the more of addicted male were found in 40 to 59 years age category and the second rank was of 18 to 39 years adult male. In comparison of high blood pressure with tobacco addiction in ≥ 40 years adult males, the results indicated that 31 per cent and 36 per cent adults were having the high systolic and diastolic blood pressure, respectively. In comparison of body mass index and tobacco addiction of ≥ 18 years adult male the data indicated that there is no effect of tobacco addiction on the BMI of adults. The information regarding the education of adults was also collected and 44 per cent females were illiterate and very less subjects were getting the graduation and post graduation degree.

CONCLUSION

The study concludes that a majority of girls had malnourishment symptoms compared to boys. More of tobacco addiction was found in adult male and 40 to 49 years adults were prone to high blood pressure problem. Illiteracy was higher in female which is indirectly responsible for the deteriorating health of the community. Adolescents are expected to enjoy good health, but this does not seem to be the case in the rural areas of developing countries like India, where poverty,

malnutrition and repeated infection are rampant. Maternal nutrition intervention programmes need to examine the role of micronutrient rich foods. Interventions to improve preconceptional maternal nutritional status for rural young girls may be more beneficial than those during pregnancy.

Paper received on : December 15, 2016

Accepted on : January 04, 2017

REFERENCES

Bulatao, R. A. and Stephens, P. W. 1992. Global estimates and projections of mortality by cause 1970-2015, Working Paper Series No. 1007 (Population and Human Resources Department, The World Bank).

Johnson, F., Wardle, J. and Griffith, J. 2002. The adolescent food habits checklist: reliability and validity of a measure of healthy eating behavior in adolescents, *European Journal of Clinical Nutrition*, 56, 644-649.

Joshi, S.B., Kanade, A.N. and Rao, S. 1998. Growth pattern of rural Indian boys aged 8-18 years: A longitudinal study, *Indian Journal of Nutrition and Dietetics*, 35, 149-156.

Kanade A.N., Joshi S.B. and Rao S. 1999. Undernutrition and adolescent growth among rural Indian boys, *Indian Pediatrics*, 36, 145-156.

Kirchengast, S. and Winkler, E.M. 1996. Nutritional status as indicator for reproductive success in Kung San and Kavango females from Namibia, *Anthropological Anz.*, 54(3), 267-276.

Naidu, A. N. and Rao, N. P. 1994. Maternal body mass index; a measure of the nutritional status in Indian populations, *European Journal of Clinical Nutrition*, 48(3), S131-S140.

National Institute of Nutrition 1991. Body mass index and mortality rates: a 10 year retrospective study, Annual Report (1989-90), Hyderabad.

NFHS III 2005. The major deficiencies are anemia, goiter, calcium deficiency and these are referred to in the Adult macro and micro malnutrition.

Rao S., Joshi S.B. and Kanade A.N. 2000b. Growth in some physical dimensions in relations to adolescent growth spurt among rural Indian boys. *Annual Human Biology*, 27, 127-138.

Rao, S., Joshi, S.B. and Kanade, A.N. 1998a. Height, weight and menarcheal age of rural Indian girls. *Indian Pediatrics*, 35, 619-626.

Rao, S., Kanade, A.N. and Joshi, S. B. 1998b. Height, weight and menarcheal age of rural Indian girls, *Indian Journal of Nutrition and Dietetics*, 35, 117-28.

Thance, M., Wilks, R.J., McFarlane, A.N., Bennett, F.I. and Forrester, T.E. 1997. Relationship between maternal nutritional status and infant's weight and body proportions at birth. *European Journal of Clinical Nutrition*, 51(3), 134-138.

WHO report, Life in the 21st century – A vision for all, Geneva.

World Bank Report 2005. Malnourishment among children in India, a regional analysis – Indian Rank.