

# Clinico-Epidemiological Study of Bovine Forestomach Disorders

Sanjay Kumar Biradar<sup>1</sup>, Raghuvir H. Bhatt<sup>2\*</sup>, Nilesh R. Padaliya<sup>1</sup>, Vaibhav D. Dodia<sup>2</sup>, Jignesh V. Vadalia<sup>1</sup>

## ABSTRACT

The present study was conducted to determine the clinico-epidemiological status of bovine forestomach disorders among 11,047 bovines which were presented during the period of three years (January 2021- December 2023) at Veterinary Clinical Complex of the College in Junagadh (Gujarat). Among these, 2279 (20.63%) were suffering from different surgical diseases, and out of them, 239 (10.48%) were diagnosed with different forestomach disorders. Age wise highest incidence was recorded in the age group of 6 to 10 years (60.66%), followed by up to 5 years (26.77%), 11 to 15 years (10.87%) and 16 to 20 years (1.67%). Females (89.53%) were more affected as compared to male animals (10.46%), and breed wise distribution showed a higher number of cases in Jaffarabadi buffaloes (43.09%), followed by non-descript buffaloes (21.75%), Gir cattle (20.50%), non-descript cattle (8.78%), Kankrej cattle (5.43%) and Holstein Friesian cattle (0.41%). The different type of forestomach disorders recorded were traumatic reticuloperitonitis (27.61%), diaphragmatic hernia (20.92%), traumatic reticulopericarditis (20.92%), rumen impaction (16.31%), traumatic reticulitis (5.43%), omasal impaction (4.18%), reticular abscess (2.92%), rumen fistula (0.83%) and rumen hernia (0.83%). The findings signify the need of early diagnosis and management of these cases to safeguard the life of animal and economy of the farmer.

**Key words:** Bovine, Diaphragmatic hernia, Incidence, Forestomach, Traumatic reticuloperitonitis.

*Ind J Vet Sci and Biotech* (2024): 10.48165/ijvsbt.20.5.32

## INTRODUCTION

The bovine forestomach including rumen, reticulum and Omasum plays a very crucial role in the production losses of bovines (Radostits *et al.*, 2006). The disorders of these compartments are characterized by similar set of clinical signs such as repeated bloat, anorexia, decreased milk yield, dullness and depression, which makes them difficult to differentiate from each other (Sharma *et al.*, 2015). These conditions primarily arise from the consumption of foreign objects. Non-digestible materials such as plastic, leather, or ropes can lead to blockages (Priyanka and Dey, 2018), while sharp metallic objects can result in inflammatory conditions of the reticulum and peritoneum. Factors that play role in ingesting these foreign bodies include indiscriminate feeding behaviour, mineral deficiency diseases, improper management techniques and urbanization (Semieka, 2010). In the times of fodder scarcity, feeding of dry feeds like hay and machine made wheat straw may lead to development of rumeno-omasal impactions (Hussain *et al.*, 2013). Development of rumen fistula may be due to traumatic injuries, while ventral herniation of rumen can be due to blunt injuries or progressive weakening of abdominal muscles (Abate, 2021). Forestomach disorders pose huge losses to dairy farmers such as production losses, economic losses and death of animals. Therefore diagnosing and managing these disorders in early stages becomes prime importance. The present study was aimed to know the retrospective status of forestomach disorders and their predisposing factors in dairy bovines.

<sup>1</sup>Department of Veterinary Surgery and Radiology, College of Veterinary Science and Animal Husbandry, Kamdhenu University, Junagadh-362 001, Gujarat, India.

<sup>2</sup>Department of Veterinary Clinical Complex, College of Veterinary Science and Animal Husbandry, Kamdhenu University, Junagadh-362 001, Gujarat, India.

**Corresponding Author:** Raghuvir H. Bhatt, Department of Veterinary Clinical Complex, College of Veterinary Science and Animal Husbandry, Kamdhenu University, Junagadh-362 001, Gujarat, India. e-mail: rhbhatt@kamdhenuuni.edu.in

**How to cite this article:** Biradar, S. K., Bhatt, R. H., Padaliya, N. R., Dodia, V. D., & Vadalia, J. V. (2024). Clinico-Epidemiological Study of Bovine Forestomach Disorders. *Ind J Vet Sci and Biotech*. 20(5), 157-159.

**Source of support:** Nil

**Conflict of interest:** None

**Submitted** 13/3/2024 **Accepted** 20/05/2024 **Published** 10/09/2024

## MATERIALS AND METHODS

The present retrospective study was carried out based on 11,047 bovine cases presented at the Veterinary Clinical Complex, College of Veterinary Science and Animal Husbandry, Kamdhenu University, Junagadh, Gujarat, India during the period of three years (January 2021 - December 2023). Out of these cases, 2279 (20.63%) animals were suffering from different surgical diseases, of which 239 (10.48 %) animals were diagnosed with different forestomach disorders. These confirmed cases were further analyzed based on age (up to 5, 6-10, 11-15 and 16-20 years), breed

(Jaffarabadi, non-descript buffalo, Gir, non-descript cattle, Kankrej and HF cattle), sex (male and female) and types of forestomach disorders.

## RESULTS AND DISCUSSION

### Age-wise Distribution of Affected Animals

In the present study, highest number of cases of forestomach disorders were recorded in the age group of 6 to 10 years (60.66%; 145/239), followed by up to 5 years (26.77%; 64/239), 11 to 15 years (10.87%; 26/239) and 16 to 20 years (1.67%; 4/239).

Higher frequency of the cases in the age group of 6 to 10 years may be attributed to the pregnancy and parturition of the animals. Descent of gravid uterus into the abdomen and its increasing size pushes rumen and reticulum, which favours the penetration of foreign bodies into wall of reticulum (Sharma *et al.*, 2015). Bouts of parturition increases the intra-abdominal pressure thereby producing more pressure on the internal organs and diaphragm leading to penetration of foreign bodies, which subsequently tear of diaphragm (Singh, 2002). Deficiency of calcium and phosphorus due to high milk production predisposes for pica (Ghurashi *et al.*, 2009). Similar findings were observed by Aref and Abdel-Hakiem (2013) in a study of sharp foreign body syndrome; Hussain *et al.* (2013) in cases of omasal impaction; Al-Abbadi *et al.* (2014) in hardware disease; Rajput *et al.* (2018) in foreign body syndrome, and Madan *et al.* (2018) in traumatic reticuloperitonitis cases.

### Sex-wise Distribution of Affected Animals

During the present study, females (89.53%; 214/239) were more affected as compared to males (10.46%; 25/239). It may be attributable to the fact that, rearing of female animals is more as compared to males. Females are reared in abundance for milk production, while very low numbers of males are kept either for breeding or agricultural purposes. Similar findings were observed by Sharma *et al.* (2015) and Lakhpati *et al.* (2019) in their studies.

### Breed-wise Distribution of Affected Animals

In the present study, breed-wise distribution showed higher incidence of forestomach disorders in buffaloes than cattle. Incidence was highest in Jaffarabadi buffaloes (43.09%)

followed by non-descript buffaloes (21.75%), Gir cattle (20.50%), non-descript cattle (8.78%), Kankrej cattle (5.43%) and Holstein Friesian cattle (0.41%) (Table 1).

**Table. 1:** Breed-wise distribution of affected animals (n=239)

Breed	No. of cases	Percentage
Jaffarabadi	103	43.09%
Non-descript buffalo	52	21.75%
Gir	49	20.50%
Non-descript cattle	21	8.78%
Kankrej	13	5.43%
Holstein Friesian	1	0.41%

Buffaloes are more affected as compared to cattle due to their pronounced indiscriminate feeding behaviour. Similar findings were noticed by Ramprabhu *et al.* (2003), Anwar *et al.* (2013), and Sharma *et al.* (2015). In the present study, breeds like Jaffarabadi, Gir and Kankrej were more affected due to their higher population in the native region of Saurashtra. Jaffarabadi and Gir are used extensively in this area for milk production and Kankrej mainly for draft purposes.

### Type of Forestomach Disorders

In the present study, higher incidence of traumatic reticuloperitonitis was observed, which was followed by diaphragmatic hernia, traumatic reticulopericarditis, rumen impaction, traumatic reticulitis, omasal impaction, reticular abscess, rumen fistula and rumen hernia (Table 2).

Higher incidence of traumatic reticuloperitonitis (TRP) can be due to their earlier presentation of the suffering animals, which if not treated may develop as diaphragmatic hernia or traumatic reticulopericarditis. Diaphragmatic hernia is commonly observed in buffaloes (Saini *et al.*, 2001) but also rarely reported in pure and cross-bred cattle (Saini *et al.*, 2007). Incidence of impaction of rumen and omasum increases with the age (Sharma *et al.*, 2015) and reticular abscess is also a complication of TRP or foreign body syndrome (Athar *et al.*, 2010), which occurs due to contamination by anaerobic microorganisms. Incidence of rumen fistula is less among animals but occurrence of rumen hernia is rare.

**Table. 2:** Type of forestomach disorders (n=239)

Disease	Cattle	Buffaloes	Total	%
Traumatic reticuloperitonitis	21	45	66	27.61
Diaphragmatic hernia	08	42	50	20.92
Traumatic reticulopericarditis	17	33	50	20.92
Rumen impaction	24	15	39	16.31
Traumatic reticulitis	6	7	13	5.43
Omasal impaction	4	6	10	4.18
Reticular abscess	0	7	7	2.92
Rumen fistula	1	1	2	0.82
Rumen hernia	0	2	2	0.82



The present study concludes that forestomach disorders are more commonly observed in the animals of age group 6-10 years and females are more affected compared to males. Among breeds Jaffarabadi recorded higher number of cases followed by non-descript buffaloes, Gir, non-descript cattle, Kankrej and HF breeds. Among different forestomach affections, incidence of traumatic reticuloperitonitis is found higher in bovines.

## ACKNOWLEDGEMENT

Authors are grateful to the authorities of Kamdhenu University and the Principal, College of Veterinary Science & AH, Junagadh for the support and facilities provided.

## REFERENCES

- Abate, M.K. (2021). Traumatic ruminal fistula in cow: A case report. *International Journal of Veterinary Sciences and Animal Husbandry*, 6(2), 23-25.
- Al-Abadi, O.S., Abu-Seida, A.M., & Al-Hussainy, S.M. (2014). Studies on rumen magnet usage to prevent hardware disease in buffaloes. *Veterinary World*, 7(6), 408-411.
- Anwar, K., Khan, I., Aslam, A., Mujtaba, M., Din, A., Amin, Y., & Ali, Z. (2013). Prevalence of indigestible rumen and reticulum foreign bodies in Achai cattle at different regions of Khyber Pakhtunkhwa. *ARPJ Journal of Agricultural and Biological Science*, 8(8), 580-586.
- Aref, N.E.M., & Abdel-Hakim, M. A. (2013). Clinical and diagnostic methods for evaluation of sharp foreign body syndrome in buffaloes. *Veterinary World*, 6(9), 586-590.
- Athar, H., Mohindroo, J., Kumar, A., Singh, K., & Sangwan, V. (2010). Diagnosis and surgical management of reticular abscess in bovines. *Indian Journal of Veterinary Surgery*, 31(1), 33-36.
- Ghurashi, M.A.H., Seri, H.I., Bakheit, A.H., & Ashwag, E.A.M. (2009). Effect of surgical removal of foreign body from goat's rumen with special reference to the prevalence of foreign body in goats in southern Darfur. *Australian Journal of Basic and Applied Sciences*, 3(2), 664-668.
- Hussain, S.A., Uppal, S.K., Randhawa, C., Sood, N.K., & Mahajan, S.K. (2013). Clinical characteristics, hematology, and biochemical analytes of primary omasal impaction in bovines. *Turkish Journal of Veterinary and Animal Sciences*, 37(3), 329-336.
- Lakhpatri, A.S., Ambore, B.N., Wankhede, G., Patodkar, V., Digraaskar, S., Suryawanshi, R., & Mali, A. (2019). Foreign body syndrome in cattle: An incidence and clinico-haemato-biochemical changes. *International Journal of Current Microbiological Applied Science*, 8(9), 783-798.
- Madan, P., Rishi, T., & Chaudhary, R.N. (2018). Ultrasonographic, radiological and surgical findings of traumatic reticuloperitonitis in buffaloes. *Haryana Veterinarian*, 57(2), 143-146.
- Priyanka, M., & Dey, S. (2018). Ruminal impaction due to plastic materials - An increasing threat to ruminants and its impact on human health in developing countries. *Veterinary World*, 11(9), 1307.
- Radostits, O.M., Gay, C., Hinchcliff, K.W., & Constable, P.D. (Eds.). (2006). *Veterinary Medicine E-Book: A textbook of the diseases of cattle, horses, sheep, pigs and goats*. Elsevier Health Sciences.
- Rajput, P.K., Parikh, P.V., Parmar, J.J., Mehta, T.A., & Patil, D.B. (2018). Studies on foreign body syndrome in bovines of Anand district of Gujarat. *Indian Journal of Animal Research*, 52(5), 744-749.
- Ramprabhu, R., Dhanapalan, P., & Prathaban, S. (2003). Comparative efficacy of diagnostic tests in the diagnosis of traumatic reticuloperitonitis and allied syndromes in cattle. *Israel Journal of Veterinary Medicine*, 58(2/3), 68-72.
- Saini, N.S., Kumar, A., Mahajan, S.K., & Sood, A.C. (2007). The use of ultrasonography, radiography, and surgery in the successful recovery from diaphragmatic hernia in a cow. *The Canadian Veterinary Journal*, 48(7), 757.
- Saini, N.S., Sobti, V.K., & Singh, S.S. (2001). Diaphragmatic hernia in cows: A study of 10 clinical cases. *Indian Journal of Veterinary Surgery*, 22, 52-53.
- Semieka, M.A. (2010). Radiography of unusual foreign body in ruminants. *Veterinary World*, 3(10), 473.
- Sharma, A.K., Dhaliwal, P.S., & Randhawa, C.S. (2015). Epidemiological studies on forestomach disorders in cattle and buffaloes. *Veterinary World*, 8(9), 1063.
- Singh, M. (2002). Evaluation of surgically treated gastrointestinal disorders in bovines. *M.V. Sc. Thesis*. Punjab Agricultural University, Ludhiana, India.